arXiv:1605.02930v2 [math.DG] 27 Jun 2016

THE CONSTANT WIDTH MEASURE SET, THE SPHERICAL MEASURE SET AND ISOPERIMETRIC EQUALITIES FOR PLANAR OVALS

MICHAŁ ZWIERZYŃSKI

ABSTRACT. In this paper we introduce and study sets: the *Constant Width Measure Set* and the *Spherical Measure Set*, which measure the constant width property and the spherical property of planar regular simple curves, respectively.

Using these sets and the Wigner caustic, we find the exact relations between the length and the area bounded by a closed regular simple convex planar curve M. Namely, the following equalities are fulfilled:

$$L_M^2 = 4\pi A_M + 8\pi \left|\widetilde{A}_{\mathrm{E}_{\frac{1}{2}}(M)}\right| + \pi \left|\widetilde{A}_{\mathrm{CWMS}(M)}\right|, \tag{0.1}$$

$$L_M^2 = 4\pi A_M + 4\pi \left|\widetilde{A}_{\mathrm{SMS}(M)}\right|, \tag{0.2}$$

where $L_M, A_M, \widetilde{A}_{\mathrm{E}_{\frac{1}{2}}(M)}, \widetilde{A}_{\mathrm{CWMS}(M)}, \widetilde{A}_{\mathrm{SMS}(M)}$ are the length of M, the area bounded by M, the oriented area of the Wigner caustic of M, the oriented area of the Constant Width Measure Set of M and the oriented area of the Spherical Measure Set of M, respectively. The equality (0.2) holds also if M is a regular closed simple curve.

We also present a new proof of Four–Vertex Theorem for generic ovals based on the number of singular points of the SMS.

1. INTRODUCTION

The *classical isoperimetric inequality* in the Euclidean plane $\mathbb{R}^2$ states that:

Theorem 1.1. *(Isoperimetric inequality) Let M be a simple closed curve of the length L_M, enclosing a region of the area A_M, then*

$$L_M^2 \geqslant 4\pi A_M, \tag{1.1}$$

and the equality (1.1) holds if and only if M is a circle.

This inequality was already known by the ancient Greeks. The first mathematical proof of this famous statement was given in the nineteenth century by Steiner [39]. After that, there have been many new proofs, generalizations, and applications of this theorem, see [3, 13, 16, 17, 22, 23, 27, 32, 39, 46], and the literature therein. In [46] recently we prove the improved isoperimetric inequality, which also gives the isoperimetric equality for convex curves of constant width.

2010 *Mathematics Subject Classification.* Primary: 52A38, 53A04, 58K70. Secondary: 52A40.

Key words and phrases. four–vertex theorem, equidistant, isoperimetric equality, isoperimetric inequality, convex curves, singularities, stability, Wigner caustic.

Theorem 1.2. *(Improved isoperimetric inequality 1) Let M be a closed regular simple convex curve. Then*

$$L_M^2 \geqslant 4\pi A_M + 8\pi \left| \widetilde{A}_{E_{\frac{1}{2}}(M)} \right|,$$

where $\widetilde{A}_{E_{\frac{1}{2}}(M)}$ is the oriented area of the Wigner caustic of M, and the equality holds if and only if M is a curve of constant width.

An affine equidistant is the set of points of chords connecting points on M where tangent lines to M are parallel, which divide the chord segments between the base points with a fixed ratio λ. If the ratio λ is equal to $\frac{1}{2}$ then this set is also known as the *Wigner caustic* of M. The Wigner caustic was first introduced by Berry, in his celebrated 1977 paper [1] on the semiclassical limit of Wigner's phase-space representation of quantum states. There are many papers considering affine equidistants and in particular the Wigner caustic, see [4, 7, 8, 9, 10, 11, 18, 20, 29, 38, 45, 46], and the literature therein. The Wigner caustic is also known as the *area evolute* ([4, 18]) or as the *symmetry defect* ([29]). Singularities of the Wigner caustic for ovals occur exactly from an antipodal pair (the tangents at the two points are parallel and the curvatures are equal), the well-known Blaschke-Süss theorem states that there are at least three pairs of antipodal points on an oval ([21, 33]). The construction of the Wigner caustic also leads to one of the two constructions of bi-dimensional improper affine spheres, which can be generelized to higher even dimensions ([5]).

Offset curves and surfaces are well-known geometric objects in the field of mathematics and computer aided geometric design, possibly because they give a powerful tool in many applications (see [12, 19, 28, 31, 36]). The Spherical Measure Set is an offset.

The classical four–vertex theorem states that the curvature function of a simple, closed, smooth plane curve has at least four local extrema. Four–Vertex Theorem was first proved for ovals in 1909 by Mukhopadhyaya ([34]) and in general by Kneser in 1912 using a projective argument ([30]), see also [6, 35] and the literature therein. We present a new proof of Four–Vertex Theorem for generic ovals and the proof is a simply consequence of the number of singular points of the Spherical Measure Set of an oval – see Corollary 4.8.

There are many important inequalities in the convex geometry and differential geometry, such as the isoperimetric inequality, the Brunn–Minkowski inequality, Aleksandrov–Fenchel inequality, Gage's inequality. The stability property of them are of great interest in geometric analysis, see [15, 16, 24, 27, 37, 40, 41] and the literature therein.

The paper is organized as follows.

In Section 2 we present geometric quantities, their Fourier series and affine equidistants, including the Wigner caustic.

Section 3 contains a definition of the Constant Width Measure Set and properties of this set for planar ovals.

In Section 4 we introduce the Spherical Measure Set and present properties of this set for planar simple regular closed curves.

Section 5 is devoted to prove the main results, isoperimetric equalities for planar simple regular closed curves.

In Section 6 we study the stability property of the isoperimetric inequality involving the Constant Width Measure Set and thanks to it we find the lower bounds of the absolute value of the oriented area of the Wigner caustic, the Constant Width Measure Set and the Spherical Measure Set.

2. Geometric quantities, affine equidistants and the Fourier series

Let M be a smooth planar curve, i.e. the image of the C^∞ smooth map from an interval to $\mathbb{R}^2$. A smooth curve is called *closed* if it is the image of a C^∞ smooth map from S^1 to $\mathbb{R}^2$. A smooth curve is called *regular* if its velocity does not vanish. A regular closed curve is called *convex* if its signed curvature has a constant sign. An *oval* is a smooth regular closed convex curve which is simple, i.e. it has no self–intersections.

Definition 2.1. Let M be a regular closed curve. A pair $a, b \in M$ $(a \neq b)$ is called the *parallel pair* if tangent lines to M at points a, b are parallel.

Definition 2.2. A *chord* passing through a pair $a, b \in M$, is the line:

$$l(a,b) = \left\{ \lambda a + (1-\lambda)b \,\middle|\, \lambda \in \mathbb{R} \right\}.$$

Definition 2.3. An affine λ-equidistant is the following set.

$$\mathrm{E}_\lambda(M) = \left\{ \lambda a + (1-\lambda)b \;\middle|\; a, b \text{ is a parallel pair of } M \right\}.$$

The set $\mathrm{E}_{\frac{1}{2}}(M)$ will be called the *Wigner caustic* of M.

Note that, for any given $\lambda \in \mathbb{R}$ we have $\mathrm{E}_\lambda(M) = \mathrm{E}_{1-\lambda}(M)$. Thus, the case $\lambda = \dfrac{1}{2}$ is special. In particular $\mathrm{E}_0(M) = \mathrm{E}_1(M) = M$.



Figure 1. An oval M and (i) $\mathrm{E}_{\frac{1}{2}}(M)$, (ii) $\mathrm{E}_{\frac{2}{5}}(M)$.

Remark 2.4. It is well known that if M is a generic oval, then for a generic λ the set $\mathrm{E}_\lambda(M)$ is a smooth closed curve with at most cusp singularities [1, 18], the number of cusps of $\mathrm{E}_{\frac{1}{2}}(M)$ is odd and not smaller than 3 [1, 18] and the number of cusps of $\mathrm{E}_\lambda(M)$ for a generic value of $\lambda \neq \dfrac{1}{2}$ is even [11].

Definition 2.5. An oval M is said to have *constant width* if the distance between every pair of parallel tangent lines to M is constant. This constant is called the *width* of the curve and we denote it by w_M.

Definition 2.6. An *average width* of an oval M is $\overline{w}_M = \dfrac{L_M}{\pi}$.

When an oval M has the constant width property, then from Barbier's theorem [2] one can get that $\overline{w}_M = w_M$.

Let us recall some basic facts about plane ovals which will be used later. The details can be found in the classical literature [23, 26].

Let M be a positively oriented oval. Take a point Θ inside M as the origin of our frame. Let p be the oriented perpendicular distance from Θ to the tangent line at a point on M, and θ the oriented angle from the positive x_1-axis to this perpendicular ray. Clearly, p is a single-valued periodic function of θ with period 2π and the parameterization of M in terms of θ and $p(\theta)$ is as follows

$$\gamma(\theta) = \big(\gamma_1(\theta), \gamma_2(\theta)\big) = \big(p(\theta)\cos\theta - p'(\theta)\sin\theta, p(\theta)\sin\theta + p'(\theta)\cos\theta\big). \tag{2.1}$$

The couple $\big(\theta, p(\theta)\big)$ is usually called the *polar tangential coordinate* on M, and $p(\theta)$ its *Minkowski support function*.

Then, the curvature κ of M is in the following form:

$$\kappa(\theta) = \frac{d\theta}{ds} = \frac{1}{p(\theta) + p''(\theta)} > 0, \tag{2.2}$$

or equivalently, the radius of a curvature ρ of M is given by:

$$\rho(\theta) = \frac{ds}{d\theta} = p(\theta) + p''(\theta).$$

Let L_M and A_M be the length of M and the area it bounds, respectively. Then one can get that

$$L_M = \int_M ds = \int_0^{2\pi} \rho(\theta) d\theta = \int_0^{2\pi} p(\theta) d\theta, \tag{2.3}$$

and

$$\begin{aligned} A_M &= \frac{1}{2}\int_M p(\theta) ds \\ &= \frac{1}{2}\int_0^{2\pi} p(\theta)\left[p(\theta) + p''(\theta)\right] d\theta = \frac{1}{2}\int_0^{2\pi} \left[p^2(\theta) - p'^2(\theta)\right] d\theta. \end{aligned} \tag{2.4}$$

The formulas (2.3) and (2.4) are known as *Cauchy's formula* and *Blaschke's formula*, respectively.

Since the Minkowski support function of M is smooth bounded and 2π–periodic, its Fourier series is in the form:

$$p(\theta) = a_0 + \sum_{n=1}^{\infty} \big(a_n \cos n\theta + b_n \sin n\theta\big). \tag{2.5}$$

Differentiation of (2.5) with respect to θ gives

$$p'(\theta) = \sum_{n=1}^{\infty} n\big(-a_n \sin n\theta + b_n \cos n\theta\big). \tag{2.6}$$

By (2.5), (2.6) and the Parseval identity one can express L_M and A_M in terms of Fourier's coefficients of $p(\theta)$ in the following way:

$$L_M = 2\pi a_0, \tag{2.7}$$

$$A_M = \pi a_0^2 - \frac{\pi}{2}\sum_{n=2}^{\infty}(n^2-1)(a_n^2+b_n^2). \tag{2.8}$$

One can notice that $\gamma(\theta), \gamma(\theta+\pi)$ is a parallel pair of M, hence γ_λ - the parameterization of $\mathrm{E}_\lambda(M)$ is as follows:

$$\begin{aligned}\gamma_\lambda(\theta) &= \big(\gamma_{\lambda,1}(\theta), \gamma_{\lambda,2}(\theta)\big)\\ &= \lambda\gamma(\theta) + (1-\lambda)\gamma(\theta+\pi)\\ &= \left(P_\lambda(\theta)\cos\theta - P'_\lambda\sin\theta, P_\lambda(\theta)\sin\theta + P'_\lambda(\theta)\cos\theta\right),\end{aligned} \tag{2.9}$$

where $P_\lambda(\theta) = \lambda p(\theta) - (1-\lambda)p(\theta+\pi)$, $\theta\in[0,2\pi]$.

Remark 2.7. If $\lambda = \dfrac{1}{2}$, then the map $M \ni \gamma(\theta) \mapsto \gamma_{\frac{1}{2}}(\theta) \in \mathrm{E}_{\frac{1}{2}}(M)$ for $\theta\in[0,2\pi]$ is the double covering of the Wigner caustic of M.

Let us also notice that in particular

$$P_{\frac{1}{2}}(\theta) = \frac{1}{2}(p(\theta) - p(\theta+\pi)) \tag{2.10}$$

$$= \sum_{\substack{n=1,\\ n \text{ is odd}}}^{\infty} \big(a_n\cos n\theta + b_n\sin n\theta\big). \tag{2.11}$$

Let $L_{\mathrm{E}_\lambda(M)}, \widetilde{A}_{E_\lambda(M)}$ denote the length of $\mathrm{E}_\lambda(M)$ and the oriented area of $\mathrm{E}_\lambda(M)$, respectively.

In [11] we show that $2L_{\frac{1}{2}}(M) \leqslant L_M$ if M is an oval.

One can check that the oriented area of the Wigner caustic has the following formula in terms of coefficients of the Fourier series of the Minkowski support function p.

$$\begin{aligned}2\widetilde{A}_{\mathrm{E}_{\frac{1}{2}}(M)} &= \frac{1}{2}\int_0^{2\pi}\Big[P^2_{\frac{1}{2}}(\theta) - P'^2_{\frac{1}{2}}(\theta)\Big]d\theta\\ &= -\frac{\pi}{2}\sum_{\substack{n=2,\\ n \text{ is odd}}}^{\infty}(n^2-1)(a_n^2+b_n^2).\end{aligned} \tag{2.12}$$

Remark 2.8. Let us notice that M is an oval of constant width if and only if coefficients a_{2n}, b_{2n} for $n\geqslant 1$ in the Fourier series of the Minkowski support function p are all equal to zero [14, 23].

3. The Constant Width Measure Set

Definition 3.1. Let M be a positively oriented oval. The *Constant Width Measure Set* of M is the following set:

$$\mathrm{CWMS}(M) = \Big\{a - b + \overline{w}_M\cdot \mathfrak{n}(a) \;\Big|\; a, b \text{ is a parallel pair of } M\Big\}, \tag{3.1}$$

where $\overline{w}_M$ is a average width of M and $\mathtt{n}(a)$ is a continuous unit normal vector field to M at a compatible with the orientation of M. We treat CWMS(M) as a subset of a vector space $V = \mathbb{R}^2$.

To visualize CWMS(M) and M in the same picture let us assume that Θ is the origin of V.

In Fig. 2 there is an example of a vector which belongs to CWMS(M) and in Fig. 3 there is an example of CWMS(M) of an oval M.



FIGURE 2. An oval M, a parallel pair a, b, tangent lines to M at a and b, vectors $a - b, \overline{w}_M \cdot \mathtt{n}(a)$ and $a - b + \overline{w}_M \cdot \mathtt{n}(a)$.

Theorem 3.2. *M is an oval of constant width if and only if CWMS$(M) = \{\Theta\}$.*

Proof. If CWMS$(M) = \{\Theta\}$, then $b - a = \overline{w}_M \cdot \mathtt{n}(a)$ for all parallel pairs a, b. This gives that the length of $b - a$ is always constant and equal to $\overline{w}_M$ and consequently M is an oval of constant width.

If M is an oval of constant width w_M, then one can notice that if a, b is a parallel pair of M, then $a - b$ is perpendicular to the tangent lines to M at a and b. This gives $a - b + w_M \cdot \mathtt{n}(a) = \Theta$. □

Theorem 3.3. *Let M be a positively oriented oval. Let $\kappa_M(a)$ denote the curvature of M at a. Let a, b be a parallel pair of M and let $q = a - b + \overline{w}_M \cdot \mathtt{n}(a)$ be a non–singular point of CWMS(M). Then*

(i) the tangent line to CWMS(M) at q is parallel to the tangent lines to M at a and b.



FIGURE 3. An oval M and CWMS(M).

(ii) the curvature of CWMS(M) at q is equal

$$\kappa_{CWMS(M)}(q) = \frac{\kappa_M(a)\cdot\kappa_M(b)}{|\kappa_M(a)+\kappa_M(b)-\overline{w}_M\cdot\kappa_M(a)\cdot\kappa_M(b)|}. \tag{3.2}$$

Proof. Let (2.1) be the parameterization of M and $p(\theta)$ its Minkowski support function.

One can check that the Minkowski support function of CWMS(M) is in the following form:

$$p_{\mathrm{CWMS}(M)}(\theta) = p(\theta)+p(\theta+\pi)-\overline{w}_M. \tag{3.3}$$

The formula (3.3) in terms of coefficients of a Fourier series of $p(\theta)$ is as follows:

$$p_{\mathrm{CWMS}(M)}(\theta) = 2\sum_{\substack{n=2,\\ n \text{ is even}}}^{\infty}(a_n\cos n\theta + b_n\sin n\theta). \tag{3.4}$$

Hence the parameterization of CWMS(M) is in the following form:

$$\begin{aligned}\gamma_{\mathrm{CWMS}(M)}(\theta) = \Big(&p_{\mathrm{CWMS}(M)}(\theta)\cos\theta - p'_{\mathrm{CWMS}(M)}(\theta)\sin\theta,\\ &p_{\mathrm{CWMS}(M)}(\theta)\sin\theta + p'_{\mathrm{CWMS}(M)}(\theta)\cos\theta\Big).\end{aligned} \tag{3.5}$$

Then:

$$\begin{aligned}\gamma'(\theta) &= \Big(p(\theta)+p''(\theta)\Big)\cdot\Big(-\sin\theta,\cos\theta\Big),\\ \gamma'_{\mathrm{CWMS}(M)}(\theta) &= \Big(p_{\mathrm{CWMS}(M)}(\theta)+p''_{\mathrm{CWMS}(M)}(\theta)\Big)\cdot\Big(-\sin\theta,\cos\theta\Big).\end{aligned}$$

Therefore the tangent line to M at $\gamma(\theta)$ is parallel to the tangent line to $\mathrm{CWMS}(M)$ at non-singular point $\gamma_{\mathrm{CWMS}(M)}(\theta)$.

By (2.5) one can get that

$$\begin{aligned}\rho_M(\theta) &= p(\theta) + p''(\theta)\\ &= a_0 - \sum_{n=1}^{\infty}(n^2-1)(a_n\cos n\theta + b_n \sin n\theta),\end{aligned} \tag{3.6}$$

$$\rho_M(\theta+\pi) = a_0 - \sum_{n=1}^{\infty}(-1)^n(n^2-1)(a_n\cos n\theta + b_n \sin n\theta). \tag{3.7}$$

By (3.4), (3.6), (3.7) one can get that

$$\begin{aligned}\rho_{\mathrm{CWMS}(M)}(\theta) &= |p_{\mathrm{CWMS}(M)}(\theta) + p''_{\mathrm{CWMS}(M)}|\\ &= \left|-2\sum_{\substack{n=2,\\ n \text{ is even}}}^{\infty}(n^2-1)(a_n\cos n\theta + b_n\sin n\theta\right|\\ &= |\rho_M(\theta)+\rho_M(\theta+\pi) - 2a_0|\\ &= |\rho_M(\theta)+\rho_M(\theta+\pi) - \overline{w}_M|\,.\end{aligned} \tag{3.8}$$

The signed curvature of $\mathrm{CWMS}(M)$ at $\gamma_{\mathrm{CWMS}(M)}(\theta)$ is in the following form:

$$\begin{aligned}\kappa_{\mathrm{CWMS}(M)}(\theta) &= \frac{1}{\rho_{\mathrm{CWMS}(M)}(\theta)}\\ &= \frac{\kappa_M(\theta)\kappa_M(\theta+\pi)}{\left|\kappa_M(\theta)+\kappa_M(\theta+\pi) - \overline{w}_M\cdot\kappa_M(\theta)\cdot\kappa_M(\theta+\pi)\right|}.\end{aligned}$$

□

By Theorem 3.3 one can get the following corollaries.

Corollary 3.4. *Let M be a positively oriented oval. Then the curvature of $CWMS(M)$ is positive on each regular connected component of $CWMS(M)$.*

Corollary 3.5. *Let M be a positively oriented oval of an average width equal to $\overline{w}_M$. Let $\mathtt{n}$ be a continuous unit normal vector field to M. Let a, b in M be a parallel pair. Let $\rho_M(a), \rho_M(b)$ denote the radius of curvature of M at a, b, respectively. Then $CWMS(M)$ is singular at the point $a - b + \overline{w}_M\cdot\mathtt{n}(a)$ if and only if*

$$\rho_M(a) + \rho_M(b) = \overline{w}_M. \tag{3.9}$$

Theorem 3.6. *Let M be a generic oval. Then $CWMS(M)$ is a connected smooth curve with cusp singularities and the number of cusps is a positive multiple of 4.*

Proof. $\mathrm{CWMS}(M)$ is a connected smooth curve because of (3.4). From Corollary 3.5 and from theory of Thom (1975) [44] one can get that generically $\mathrm{CWMS}(M)$ has cusp singularity when (3.9) holds. By (3.8) the condition (3.9) can be written in terms of the Fourier series of the Minkowski support function of M in the following way:

$$\rho_{\mathrm{CWMS}(M)}(\theta) = 0 \iff \sum_{\substack{n=2,\\ n \text{ is even}}}^{\infty}(n^2-1)(a_n\cos n\theta + b_n\sin n\theta) = 0. \tag{3.10}$$

Therefore the number of cusps of CWMS(M) is the number of zeros of (3.10) in the range 0 to 2π. Since $\rho_{\mathrm{CWMS}(M)}(0) = \rho_{\mathrm{CWMS}(M)}(\pi)$, the number of cusps in the range from 0 to π must be even. The function $\rho_{\mathrm{CWMS}(M)}(\theta)$ is π-periodic, hence the number of cusps in the range from 0 to 2π is a multiple of 4. Moreover, the number of cusps must be at least 4, since the Fourier series in (3.10) begins with terms in $\cos 2\theta$ and $\sin 2\theta$. □

Definition 3.7. The *tangent line to the* CWMS(M) *at a cusp point* p is the limit of a sequence of $T_{q_n}M$ for any sequence q_n of regular points of CWMS(M) converging to p.

It is easy to see that this definition does not depend on the choice of a converging sequence of regular points. From Theorem 3.3(i) we can see that the tangent line to CWMS(M) at the cusp point $a - b + \overline{w}_M \cdot \mathtt{n}(a)$ is parallel to the tangent lines to M at a and b.

It is easy to see that if M is a generic oval, then for any line l in $\mathbb{R}^2$ there exists exactly two points $a, b \in M$ in which tangent lines to M at a and b are parallel to l. Therefore we get the following corollary.

Corollary 3.8. *Let M be a generic oval. Then for any line l in $\mathbb{R}^2$ there exists exactly two points $p, q \in$ CWMS(M) such that tangent lines to CWMS(M) at p and q are parallel to l.*

We can define the normal vector to CWMS(M) at any its point $a-b+\overline{w}_M \cdot \mathtt{n}(a)$ in the following way.

Definition 3.9. The normal vector to CWMS(M) at $a - b + \overline{w}_M \cdot \mathtt{n}(a)$ is $\mathtt{n}(a)$.

Let us notice that the continuous normal vector field to CWMS(M) at regular and cusp points is perpendicular to the tangent line to CWMS(M). Using this fact and the above definition we define the rotation number in the following way.

Definition 3.10. The *rotation number* of a smooth closed curve M with at most cusp singularities is the number of rotation of its continuous normal vector field.

Proposition 3.11. *Let M be a generic oval. Then the absolute value of the rotation number of CWMS(M) is equal to one.*

Proof. It is a consequence of Theorem 3.6 and Corollary 3.8. □

Proposition 3.12. *If M is an oval, then CWMS(M) has the center of symmetry.*

Proof. Let a, b be a parallel pair of M, then $p = a - b + \overline{w}_M \cdot \mathtt{n}(a) \in \mathrm{CWMS}(M)$ and $q = b - a + \overline{w}_M \cdot \mathtt{n}(b) \in \mathrm{CWMS}(M)$, and because $\mathtt{n}(a) = -\mathtt{n}(b)$ one can notice that Θ is the center of symmetry of CWMS(M). □

Proposition 3.13. *There exists oval M for which the Constant Width Measure Set has exactly $4n$ cusp singularities.*

Proof. One can check that $p_{4n}(\theta) = 4n^2 + 2 + \cos 2n\theta$ is a Minkowski support function of an oval M_{4n} and CWMS(M_{4n}) has exactly $4n$ cusps. □

In Fig. 4 there is an example of an oval M for which the Constant Width Measure Set has exactly 12 cusp singularities.

Theorem 3.14. *Let M be an oval. Then*

$$L_{CWMS(M)} \leqslant 4L_M.$$

Proof. Let $p(\theta)$ be the Minkowski support function of M and let $p_{\mathrm{CWMS}(M)}$ (like in (3.3)) be the support function of $\mathrm{CWMS}(M)$. Then

$$\begin{aligned} L_{\mathrm{CWMS}} &= \int_0^{2\pi} |\gamma'_{\mathrm{CWMS}(M)}(\theta)|d\theta \\ &= \int_0^{2\pi} |p_{\mathrm{CWMS}(M)}(\theta) + p''_{\mathrm{CWMS}(M)}(\theta)|d\theta \\ &= \int_0^{2\pi} |p(\theta) + p''(\theta) + p(\theta+\pi) + p''(\theta+\pi) - \overline{w}|d\theta \\ &\leqslant \int_0^{2\pi} |p(\theta) + p''(\theta)|d\theta + \int_0^{2\pi} |p(\theta+\pi) + p''(\theta+\pi)|d\theta + \int_0^{2\pi} |\overline{w}|d\theta \\ &= L_M + L_M + 2L_M = 4L_M. \end{aligned}$$

□



FIGURE 4. An oval M_{12} and $\mathrm{CWMS}(M_{12})$ which has 12 cusp singularities. The support function of M_{12} is $p_{12}(\theta) = 38 + \cos 6\theta$.

4. The Spherical Measure Set

We recall definition of an offset (this set is also known as a *parallel* set).

Definition 4.1. Let M be a regular positively oriented simply closed curve. An *α-offset* of M is the following set

$$\mathrm{F}_\alpha(M) = \big\{a + \alpha \cdot \mathtt{n}(a) \;\big|\; a \in M\big\},$$

where $\mathtt{n}(a)$ is a continuous unit normal vector field to M at a compatible with the orientation of M.

It is well known that offsets generically admit at most cusp singularities, singularities of all offsets of M form an evolute of M and set of all points of self–intersections of offsets forms a medial axis of M. See [12, 19, 28, 31, 36] and the literature therein.

Definition 4.2. The *Spherical Measure Set* of a regular positively oriented simply closed curve M is an offset at level $\dfrac{L_M}{2\pi}$,

$$\mathrm{SMS}(M) = \mathrm{F}_{\frac{L_M}{2\pi}}(M) = \left\{ a + \frac{L_M}{2\pi} \cdot \mathtt{n}(a) \;\middle|\; a \in M \right\}, \tag{4.1}$$

where $\mathtt{n}$ is the unit normal vector field compatible with the orientation of M.

Remark 4.3. If M is an oval, then $\dfrac{L_M}{2\pi} = \dfrac{\overline{w}_M}{2}$.

See Fig. 5 and Fig. 6 for examples of SMS.



FIGURE 5. An oval M and $\mathrm{SMS}(M)$.



FIGURE 6. A regular simple closed curve M and $\mathrm{SMS}(M)$.

Proposition 4.4. *Let M be a regular positively oriented simply closed curve. Then SMS(M) $= \{x\}$ if and only if M is a circle and x is its center.*

Proof. If M is a circle and x is its center, then it easy to see that $\frac{1}{2}\overline{w}$ is a radius of M and $\mathrm{SMS}(M) = \{x\}$.

Let $w = \dfrac{L_M}{\pi}$. Let us assume that $\mathrm{SMS}(M) = \{x\}$. Let a, b be a parallel pair of M, then $w \cdot \mathfrak{n}(a) = 2x - 2a$ and $w \cdot \mathfrak{n}(b) = 2x - 2b$. If $\mathfrak{n}(a) = \mathfrak{n}(b)$, then $a = b$, otherwise if $\mathfrak{n}(a) = -\mathfrak{n}(b)$, then $|a - b| = w$ and $a + b = 2x$, so M is a curve of constant width and x is the center of symmetry of M, hence M is a circle. □

Proposition 4.5. *Let M be a positively oriented regular closed curve. Let κ_M denote the curvature of M at a. Then*

(i) a point $a + \dfrac{L_M}{2\pi} \cdot \mathfrak{n}(a)$ is a singular point of SMS if and only if

$$L_M \kappa_M(a) = 2\pi, \tag{4.2}$$

(ii) the tangent line to SMS(M) at non-singular point $a + \dfrac{L_M}{2\pi} \cdot \mathfrak{n}(a)$ is parallel to the tangent line to M at a point a.

(iii) the curvature of SMS(M) at non-singular point $q = a + \dfrac{L_M}{2\pi} \cdot \mathfrak{n}(a)$ is equal

$$\kappa_{SMS(M)}(q) = \frac{2\pi\kappa_M(a)}{|2\pi - L_M\kappa_M(a)|}. \tag{4.3}$$

Proof. Let $\gamma(s)$ be the arc length parameterization of M and let $\mathfrak{t}, \mathfrak{b}$ be its Frenet frame. Then a parameterization of $\mathrm{SMS}(M)$ is as follows:

$$\gamma_{\mathrm{SMS}(M)}(s) = \gamma(s) + \frac{L_M}{2\pi}\mathfrak{n}(s).$$

Then:

$$\begin{aligned}\gamma'_{\mathrm{SMS}(M)}(s) &= \mathfrak{t}(s) - \frac{L_M}{2\pi} \cdot \kappa_M(s)\mathfrak{t}(s) \\ &= \frac{1}{2\pi}\left(2\pi - L_M\kappa_M(s)\right)\mathfrak{t}(s).\end{aligned}$$

Therefore *(i), (ii)* hold.

The curvature of $\mathrm{SMS}(M)$ at non–singular $\gamma_{\mathrm{SMS}(M)}(s)$ is equal to

$$\begin{aligned}\kappa_{\mathrm{SMS}(M)}(s) &= \frac{\det\left(\gamma'_{\mathrm{SMS}(M)}(s), \gamma''_{\mathrm{SMS}(M)}(s)\right)}{\left|\gamma'_{\mathrm{SMS}(M)}(s)\right|^3} \\ &= \frac{2\pi\kappa_M(s)}{\left|2\pi - L_M\kappa_M(s)\right|}.\end{aligned}$$

□

Theorem 4.6. *Let M be a generic regular closed curve. Then the number of cusp singularities of SMS(M) is even and not smaller than 2. If M is a generic oval, then the number of cusp singularities of SMS(M) is not smaller than 4.*

Proof. Let us assume that M is positively oriented.

By theory of Thom (1975) [44] one can get that generically $\mathrm{SMS}(M)$ has cusp singularity when (4.2) holds.



FIGURE 7. The cusp singularity with a continuous normal vector field. Vectors in upper regular component of a curve are directed outside the cusp, others are directed inside the cusp.

A continuous normal vector field to the germ of a curve with the cusp singularity is directed outside the cusp on the one of two connected regular components and is directed inside the cusp on the other component as it is showed in Fig. 7.

Without loss of generality, let us assume that M is positively oriented. It is easy to see that the rotation number of $\mathrm{SMS}(M)$ is equal to the rotation number of M, which is an integer. Thus, the number of cusps of $\mathrm{SMS}(M)$ is even.

Let us assume that $\mathrm{SMS}(M)$ is regular. Let $[0, L_M) \ni s \mapsto \gamma(s) \in \mathbb{R}^2$ be the arc length parameterization of M. Then by Proposition 4.5(i) we get that the curvature at each point of M is smaller or greather than $\dfrac{2\pi}{L_M}$. If $\kappa_M(s) < \dfrac{2\pi}{L_M}$ for all s, then

$$\int_0^{L_M} \kappa_M(s)ds < \int_0^{L_M} \frac{2\pi}{L_M} ds,$$
$$2\pi < 2\pi.$$

We get the same result if we assume that $\kappa_M(s) > \dfrac{2\pi}{L_M}$ for all s, therefore $\mathrm{SMS}(M)$ got at least one cusp singularity.

Let M be a generic oval.

Let (2.1) be the parameterization of M and $p(\theta)$ its Minkowski support function.

By Remark 2.4 there are at least 3 cusps of the Wigner caustic of M, then there exist $\theta_1 < \theta_2 < \theta_3$ such that $\rho_M(\theta_i) = \rho_M(\theta_i + \pi)$ for $i = 1, 2, 3$ and by Corollary 3.5 there are at least 4 cusps of the Constant Width Measure Set of M, then there exists $\varphi_1 < \varphi_2$ such that $\rho_M(\varphi_j) + \rho_M(\varphi_j + \pi) = \overline{w}_M$ for $j = 1, 2$, therefore it is easy to see that there are at least four values of $\theta \in S^1$ such that $\rho_M(\theta) = \dfrac{1}{2}\overline{w}_M$, hence there are at least 4 cusps of the Spherical Measure Set of M. □

Remark 4.7. The minimal number of cusp singularities of a $\mathrm{SMS}(M)$ of a generic regular closed simple curve M cannot be 4 - see an example in Fig. 8.

By Theorem 4.6 we get another proof of the classical Four–Vertex Theorem for ovals.

Corollary 4.8. *Let M be a generic oval. Then M has at least four vertices.*

Proof. It is a consequence of Roll's theorem and the fact that there are at least 4 points on M in which the curvature is equal to $\dfrac{2\pi}{L_M}$ - see Proposition 4.5(i) and Theorem 4.6. □

Proposition 4.9. *If M is an oval of constant width, then $SMS(M) = E_{\frac{1}{2}}(M)$. Thus, the map $M \ni \gamma(\theta) \mapsto \gamma_{SMS(M)}(\theta) \in SMS(M)$ is the double covering of $SMS(M)$.*

If M is a curve with the center of symmetry, then $CWMS(M)$ and $SMS(M)$ are similar curves and the ratio of symmetry between them is equal to 2.

Proof. Let p be the Minkowski support function of M. Then one can check that the Minkowski support function of $\mathrm{SMS}(M)$ is in the form:

$$p_{\mathrm{SMS}(M)}(\theta) = p(\theta) - \frac{1}{2}\overline{w}_M. \tag{4.4}$$

In terms of coefficients of the Fourier series of $p(\theta)$:

$$p_{\mathrm{SMS}(M)}(\theta) = \sum_{n=1}^{\infty}(a_n \cos n\theta + b_n \sin n\theta). \tag{4.5}$$

Then Proposition 4.9 is a consequence of the support functions of $\mathrm{E}_{\frac{1}{2}}(M)$, $\mathrm{CWMS}(M)$, $\mathrm{SMS}(M)$ in terms of coefficients of the Fourier series of $p(\theta)$ - see (2.11), (3.4), (4.5).

□



FIGURE 8. A simple regular closed curve M and $\mathrm{SMS}(M)$ which has 2 cusp singularities.

Corollary 4.10. *Let M be an oval such that SMS(M) is a curve with at most cusp singularities. If M is not a curve of constant width, then the absolute value of the rotation number of SMS(M) is equal to one, otherwise it is equal to* $\frac{1}{2}$.

Proposition 4.11. *There exists oval M for which the Constant Width Measure Set has exactly n cusp singularities for $n \geqslant 3$.*

Proof. Let $n \geqslant 3$ and let

$$p_n(\theta) = \begin{cases} n^2 + 2 + \cos n\theta & \text{if} \quad n \equiv 1 (\text{mod } 2) \\ \left(\frac{n}{2}\right)^2 + 2 + \cos \frac{n\theta}{2} & \text{if} \quad n \equiv 0 (\text{mod } 4) \\ n^2 + 2 + \cos \frac{(n-2)\theta}{2} + \cos \frac{n\theta}{2} & \text{if} \quad n \equiv 2 (\text{mod } 4) \end{cases}$$

be a Minkowski support function of an oval M_n. Then one can check that SMS(M_n) has exactly n cusp singularities. □

In Fig. 9 there is an example of an oval M for which SMS(M) has exactly 10 cusp singularities.



FIGURE 9. An oval M_{10} and SMS(M_{10}) which has 10 cusp singularities. The support function of M_{10} is $p_{10}(\theta) = 102 + \cos 4\theta + \cos 5\theta$.

Theorem 4.12. *Let M be an oval. Then*

$$L_{SMS(M)} \leqslant 2L_M \tag{4.6}$$

and if M is an oval of constant width, then

$$L_{SMS(M)} \leqslant L_M \tag{4.7}$$

Proof. Let $p(\theta)$ be the Minkowski support function of M and let $p_{\text{SMS}(M)}$ (like in (4.4)) be the support function of SMS(M). Let M is not an oval of constant width.

Then

$$
\begin{aligned}
L_{\mathrm{SMS}} &= \int_0^{2\pi} |\gamma'_{\mathrm{SMS}(M)}(\theta)| d\theta \\
&= \int_0^{2\pi} |p_{\mathrm{SMS}(M)}(\theta) + p''_{\mathrm{SMS}(M)}(\theta)| d\theta \\
&= \int_0^{2\pi} \left| p(\theta) - \frac{1}{2}\overline{w}_M + p''(\theta) \right| d\theta \\
&\leqslant \int_0^{2\pi} |p(\theta) + p''(\theta)| \, d\theta + \frac{1}{2}\int_0^{2\pi} \overline{w}_M d\theta \\
&= L_M + L_M = 2L_M.
\end{aligned} \tag{4.8}
$$

By (4.8) and Proposition 4.9 the inequality (4.7) holds. □

5. Isoperimetric equalities

Theorem 5.1. *(Isoperimetric equality 1) Let M be a closed regular simple convex planar curve. Then*

$$
L_M^2 = 4\pi A_M + 8\pi \left| \widetilde{A}_{E_{\frac{1}{2}}(M)} \right| + \pi \left| \widetilde{A}_{CWMS(M)} \right|, \tag{5.1}
$$

where $L_M, A_M, \widetilde{A}_{E_{\frac{1}{2}}(M)}, \widetilde{A}_{CWMS(M)}$ are the length of M, the area bounded by M, the oriented area of the Wigner caustic of M and the oriented area of the Constant Width Measure Set of M, respectively.

Proof. Let $p(\theta)$ be the Minkowski support function of M and let (2.5) be its Fourier series. Then (3.4) is the Fourier series of the Minkowski support function $p_{\mathrm{CWMS}(M)}(\theta)$ of $\mathrm{CWMS}(M)$.

The oriented area of $\mathrm{CWMS}(M)$ in terms of coefficients of the Fourier series of p is equal

$$
\begin{aligned}
\widetilde{A}_{\mathrm{CWMS}(M)} &= \frac{1}{2}\int_0^{2\pi} \left(p^2_{\mathrm{CWMS}(M)}(\theta) - p'^2_{\mathrm{CWMS}(M)}(\theta) \right) d\theta \\
&= -2\pi \sum_{\substack{n=2, \\ n \text{ is even}}}^{\infty} (n^2-1)(a_n^2 + b_n^2).
\end{aligned} \tag{5.2}
$$

Then by (2.7), (2.8), (2.12), (5.2) one can easily verify that the equality (6.9) holds. □

Theorem 5.2. *(Isoperimetric equality 2) Let M be a closed simple regular planar curve. If M is not a curve of constant width, then*

$$
L_M^2 = 4\pi A_M + 4\pi \left| \widetilde{A}_{SMS(M)} \right|, \tag{5.3}
$$

otherwise

$$
L_M^2 = 4\pi A_M + 8\pi \left| \widetilde{A}_{SMS(M)} \right|, \tag{5.4}
$$

where $L_M, A_M, \widetilde{A}_{SMS(M)}$ are the length of M, the area bounded by M and the oriented area of the Spherical Measure Set of M, respectively.

Proof. Let M be not a curve of constant width.

Let $[0, L_M) \ni s \mapsto \gamma(s) \in M$ be an arc length parameterization of M. Then $[0, L_M) \ni s \mapsto \gamma_{\mathrm{SMS}(M)}(s) = \gamma(s) + \dfrac{L_M}{2\pi} \cdot \mathbb{n}(s) \in \mathrm{SMS}(M)$ is a parameterization of M.

Let $\mathbb{t}, \mathbb{n}, \mathbb{b}$ be the Frenet frame of M.

By Green's theorem the oriented area of $\mathrm{SMS}(M)$ is equal to

$$\begin{aligned}
\widetilde{A}_{\mathrm{SMS}(M)} &= \frac{1}{2}\int_0^{L_M} \left(\gamma_{\mathrm{SMS}(M)}(s) \times \gamma'_{\mathrm{SMS}(M)}(s)\right) \cdot \mathbb{b}(s) ds \\
&= \frac{1}{2}\int_0^{L_M} \left(\gamma(s) + \frac{L_M}{2\pi}\mathbb{n}(s)\right) \times \left(1 - \frac{L_M}{2\pi}\kappa(s)\right) \mathbb{t}(s) \cdot \mathbb{b}(s) ds \\
&= \frac{1}{2}\int_0^{L_M} \gamma(s) \times \mathbb{t}(s) \cdot \mathbb{b}(s) ds - \frac{L_M}{2\pi}\int_0^{L_M} \mathbb{b}(s) \cdot \mathbb{b}(s) ds + \\
&\quad + \frac{1}{2} \cdot \frac{L_M^2}{4\pi^2}\int_0^{L_M} \kappa(s) ds \\
&= A_M - \frac{L_M}{2\pi} \cdot L_M + \pi \cdot \frac{L_M^2}{4\pi^2} = A_M - \frac{L_M^2}{4\pi}
\end{aligned} \tag{5.5}$$

□

From the isoperimetric equalities one can get following corollaries.

Corollary 5.3. [46] *A closed regular simple convex planar curve M is a curve of constant width if and only if*

$$L_M^2 = 4\pi A_M + 8\pi \left|\widetilde{A}_{E_{\frac{1}{2}}(M)}\right|,$$

where $L_M, A_M, \widetilde{A}_{E_{\frac{1}{2}}(M)}$ are the length of M, the area bounded by M and the oriented area of the Wigner caustic of M, respectively.

Corollary 5.4. *A closed regular simple convex planar curve M is a curve which has the center of symmetry if and only if*

$$L_M^2 = 4\pi A_M + \pi \left|\widetilde{A}_{CWMS(M)}\right|,$$

where $L_M, A_M, \widetilde{A}_{CWMS(M)}$ are the length of M, the area bounded by M and the oriented area of the Constant Width Measure Set of M, respectively.

Corollary 5.5. *Let M be a closed simple convex planar curve. If M is not a curve of constant width then*

$$4\left|\widetilde{A}_{SMS(M)}\right| = 8\left|\widetilde{A}_{E_{\frac{1}{2}}(M)}\right| + \left|\widetilde{A}_{CWMS(M)}\right|, \tag{5.6}$$

otherwise

$$SMS(M) = E_{\frac{1}{2}}(M), CWMS(M) = \{\Theta\}, \tag{5.7}$$

where $\widetilde{A}_{E_{\frac{1}{2}}(M)}, \widetilde{A}_{CWMS(M)}, \widetilde{A}_{SMS(M)}$ are the oriented area of the Wigner caustic of M, the oriented area of the Constant Width Measure Set of M and the oriented area of the Spherical Measure Set of M, respectively.

Proposition 5.6. *Let M be a closed simple convex planar curve. If M is not a curve of constant width then*

$$L_{SMS(M)} \leqslant \frac{1}{2} L_{CWMS(M)} + 2L_{E_{\frac{1}{2}}(M)}. \tag{5.8}$$

Proof. By (2.10), (3.3), (4.4) and Remark 2.7 we get the following:

$$\begin{aligned}
L_{\mathrm{SMS}(M)} &= \int_0^{2\pi} \left| p_{\mathrm{SMS}(M)}(\theta) \right| d\theta \\
&= \int_0^{2\pi} \left| p(\theta) - \frac{1}{2}\overline{w}_M \right| d\theta \\
&= \frac{1}{2} \int_0^{2\pi} \left| p(\theta) + p(\theta+\pi) - \overline{w}_M + p(\theta) - p(\theta+\pi) \right| d\theta \\
&= \frac{1}{2} \int_0^{2\pi} \left| p_{\mathrm{CWMS}(M)}(\theta) + 2p_{\mathrm{E}_{\frac{1}{2}}(M)}(\theta) \right| d\theta \\
&\leqslant \frac{1}{2} \int_0^{2\pi} \left| p_{\mathrm{CWMS}(M)}(\theta) \right| d\theta + \int_0^{2\pi} \left| p_{\mathrm{E}_{\frac{1}{2}}(M)}(\theta) \right| d\theta \\
&= \frac{1}{2} L_{\mathrm{CWMS}(M)} + 2L_{\mathrm{E}_{\frac{1}{2}}(M)}
\end{aligned}$$

□

Proposition 5.7. *Let M be a generic regular simple closed curve. Then the orientation of SMS(M) is reversed against the orientation of M. If M is a generic oval, then the orientation of CWMS(M) is reversed against the orientation of M.*

Proof. Let M be positively oriented. Then the negatively orientation of $\mathrm{CWMS}(M)$ and $\mathrm{SMS}(M)$ is a consequence of (5.2) and (5.5). □

In Fig. 10 there is an example of an oval M for which

$$p(\theta) = 115 + 10\cos 2\theta + \frac{1}{3}\cos 3\theta + \sin 4\theta - 3\sin 5\theta$$

is its Minkowski support function. Then $\mathrm{E}_{\frac{1}{2}}(M)$ has 5 cusp singularities, $\mathrm{CWMS}(M)$ has 4 cusp singularities, $\mathrm{SMS}(M)$ has 10 cusp singularities and their support functions are as follows:

$$\begin{aligned}
p_{\mathrm{E}_{\frac{1}{2}}(M)}(\theta) &= \frac{1}{3}\cos 3\theta - 3\sin 5\theta, \\
p_{\mathrm{CWMS}(M)}(\theta) &= 10\cos 2\theta + \sin 4\theta, \\
p_{\mathrm{SMS}(M)}(\theta) &= 10\cos 2\theta + \frac{1}{3}\cos 3\theta + \sin 4\theta - 3\sin 5\theta.
\end{aligned}$$

6. The stability of improved isoperimetric inequalities

An n – *dimensional convex body* is a bounded convex subset in $\mathbb{R}^n$ which is closed and has interior points. Let $\mathcal{C}^n$ denote the set of all n – dimensional convex bodies.

An inequality in the convex geometry can be written as

$$\Phi(K) \geqslant 0, \tag{6.1}$$



FIGURE 10. An oval M and $\mathrm{E}_{\frac{1}{2}}(M)$ (dashed line, 5 cusps), CWMS(M) (bold line, 4 cusps), SMS(M) (normal line, 10 cusps). The Minkowski support function of M is equal to $p(\theta) = 115 + 10\cos 2\theta + \frac{1}{3}\cos 3\theta + \sin 4\theta - 3\sin 5\theta$.

where $\Phi : \mathcal{C}^n \to \mathbb{R}$ is a function and the inequality (6.1) holds for all K in $\mathcal{C}^n$. Let $\mathcal{C}^n_\Phi$ be a subset of $\mathcal{C}^n$ for which equality in (6.1) holds.

Let $L_{\partial K}$ denote the length of the boundary of K, let $A_{\partial K}$ denote the area enclosed by ∂K (i.e. the area of K).

For example, let $n = 2$ and let $\Phi(K) = L^2_{\partial K} - 4\pi A_{\partial K}$. Then the inequality $\Phi(K) \geqslant 0$ is the classical isoperimetric inequality in $\mathbb{R}^2$. In this case $\mathcal{C}^2_\Phi$ is a set of disks.

In this section we will study stability properties associated with (6.1). We ask if K must by close to a member of $\mathcal{C}^n_\Phi$ whenever $\Phi(K)$ is close to zero. Let $d : \mathcal{C}^n \times \mathcal{C}^n \to \mathbb{R}$ denoted in some sense the deviation between two convex bodies. d should satisfy two following conditions:

(i) $d(K, L) \geqslant 0$ for all $K, L \in \mathcal{C}^n$,
(ii) $d(K, L) = 0$ if and only if $K = L$.

If $\Phi, \mathcal{C}^n_\Phi$ and d are given, then the *stability problem* associated with (6.1) is as follows.

Find positive constants c, α such that for each $K \in \mathcal{C}^n$, there exists $N \in C^n_\Phi$ such that

$$\Phi(K) \geqslant c g^\alpha(K, N). \tag{6.2}$$

Let $p_{\partial K}$ and $p_{\partial N}$ be support functions of convex bodies K and N, respectively. Usually to measure the deviation between K and N one can use the *Hausdorff distance*,

$$d_\infty(K, N) = \max_\theta \Big| p_{\partial K}(\theta) - p_{\partial N}(\theta) \Big|. \tag{6.3}$$

Another such measure is the measure that corresponds to the L_2-metric in the function space. It is defined by

$$d_2(K, N) = \left(\int_0^{2\pi} \Big| p_{\partial K}(\theta) - p_{\partial N}(\theta) \Big|^2 d\theta \right)^{\frac{1}{2}}. \tag{6.4}$$

It is easy to see that $d_\infty(K, N) = 0$ (or $d_2(K, N) = 0$) if and only of $K = N$.

Definition 6.1. [46] Let p_M be the Minkowski support function of a positively oriented oval M of length L_M. Then

$$p_{W_M}(\theta) = \frac{L_M}{2\pi} + \frac{p_M(\theta) - p_M(\theta + \pi)}{2} \tag{6.5}$$

will be the support function of a curve W_M which will be called the *Wigner caustic type curve associated with M*.

Proposition 6.2. [46] *Let W_M be the Wigner caustic type curve associated with an oval M. Then W_M has the following properties:*

(i) W_M is an oval of constant width,
(ii) $L_{W_M} = L_M$,
(iii) $E_{\frac{1}{2}}(W_M) = E_{\frac{1}{2}}(M)$,
(iv) $A_{W_M} \geqslant A_M$ and the equality holds if and only if M is a curve of constant width,
(v) $W_M = M$ if and only if M is a curve of constant width.

In [46] we show the stability properties of Theorem 1.2.

Theorem 6.3. [46] *Let K be strictly convex domain of area $A_{\partial K}$ and perimeter $L_{\partial K}$ and let $\widetilde{A}_{E_{\frac{1}{2}}(\partial K)}$ denote the oriented area of the Wigner caustic of ∂K. Let W_K denote the convex body for which ∂W_K is the Wigner caustic type curve associated with ∂K. Then*

$$L^2_{\partial K} - 4\pi A_{\partial K} - 8\left|\widetilde{A}_{E_{\frac{1}{2}}(\partial K)}\right| \geqslant 4\pi^2 d^2_\infty(K, W_K), \tag{6.6}$$

where equality holds if and only if ∂K is a curve of constant width.

Theorem 6.4. [46] *Under the same assumptions of Theorem 6.3, one gets*

$$L^2_{\partial K} - 4\pi A_{\partial K} - 8\pi\left|\widetilde{A}_{E_{\frac{1}{2}}(\partial K)}\right| \geqslant 6\pi d^2_2(K, W_K), \tag{6.7}$$

where equality holds if and only if ∂K is a curve of constant width, or the Minkowski support function of ∂K is in the form

$$p_{\partial K}(\theta) = a_0 + a_2\cos 2\theta + b_2 \sin 2\theta + \sum_{\substack{n=1,\\ n\ is\ odd}}^{\infty} (a_n \cos n\theta + b_n \sin n\theta).$$

From Theorem 6.3, Theorem 6.8 and 5.1 we can get the following result.

Corollary 6.5. *Under the same assumptions of Theorem 6.3, one gets*

$$\left|\widetilde{A}_{CWMS(\partial K)}\right| \geqslant \max\left\{4\pi d^2_\infty(K, W_K),\ 6d^2_2(K, W_K)\right\}. \tag{6.8}$$

From Theorem 5.1 we can easily obtain the second improved isoperimetric inequality (see Theorem 6.6), study its stability property and obtain similar results like these in Corollary 6.5, but for the Wigner caustic of an oval.

Theorem 6.6. *(Improved isoperimetric inequality 2) Let M be a closed regular simple convex planar curve. Then*

$$L^2_M \geqslant 4\pi A_M + \pi\left|\widetilde{A}_{CWMS(M)}\right|, \tag{6.9}$$

where $L_M, A_M, \widetilde{A}_{CWMS(M)}$ are the length of M, the area bounded by M and the oriented area of the Constant Width Measure Set of M, respectively. The equality in (6.9) holds if and only if M is a centrally symmetric.

From this point let us assume that $n = 2$ and by Theorem 6.6 let

$$\Phi(K) = L^2_{\partial K} - 4\pi A_{\partial K} - \pi\left|\widetilde{A}_{\mathrm{CWMS}(\partial K)}\right|. \tag{6.10}$$

From Theorem 6.6 one can see that C^2_Φ consists of centrally symmetric bodies.

Let K be the convex body in $\mathbb{R}^2$. Let $\vec{u}$ be a unit vector in $\mathbb{R}^2$. Then the *Steiner point* which can be defined in terms of the Minkowski support function of ∂K is as follows:

$$\vec{s}(K) = \frac{1}{\pi}\int_0^{2\pi} \vec{u}(\theta)\cdot p_{\partial K}(\theta)d\theta. \tag{6.11}$$

In terms of coefficients of the Fourier series of $p_{\partial K}$ one can notice that $\vec{s}(K) = (a_1, b_1)$.

For more details on the Steiner point of a convex body see [25, 41, 42].

Definition 6.7. Let K be the convex body. Then $S_K = \frac{1}{2}(K + (-K))$ is centrally symmetric convex body, where $+$ is the Minkowski addition and $-K = \{-k \mid k \in K\}$. Often S_K is called the *Steiner symmetral of* K, and the process of generating the set from K is known as *symmetrization*. Let as translate S_K such that the center of S_K becomes the Steiner point of K.

The support function of ∂S_K is equal to

$$p_{\partial S_K}(\theta) = \vec{s}(K) + \frac{p_{\partial K}(\theta) + p_{\partial K}(\theta + \pi)}{2} \tag{6.12}$$

and in terms of coeeficients of the Fourier series of $p_{\partial K}$:

$$p_{\partial S_K}(\theta) = a_0 + a_1 \cos\theta + b_1 \sin\theta + \sum_{\substack{n=2,\\ n \text{ is even}}}^{\infty} (a_n \cos n\theta + b_n \sin n\theta)\,. \tag{6.13}$$

Theorem 6.8. *Let K be strictly convex domain of the area $A_{\partial K}$ and the perimeter $L_{\partial K}$ and let $\widetilde{A}_{CWMS(\partial K)}$ denote the oriented area of the Constant Width Measure Set of ∂K. Let S_K be the Steiner symmetral of K. Then*

$$L^2_{\partial K} - 4\pi A_{\partial K} - \pi \left|\widetilde{A}_{CWMS(\partial K)}\right| \geqslant 8\pi^2 d^2_\infty(K, S_K), \tag{6.14}$$

where equality holds if and only if ∂K is centrally symmetric.

Proof. By (2.5), (2.6), (2.3), (2.4), one can get the Fourier series of Φ (see (6.10)):

$$\begin{aligned} \Phi(K) &= L^2_{\partial K} - 4\pi A_{\partial K} - 8\pi \left|\widetilde{A}_{E_{\frac{1}{2}}(M)}\right| \\ &= 2\pi^2 \sum_{\substack{n=3,\\ n \text{ is odd}}}^{\infty} (n^2 - 1)(a_n^2 + b_n^2). \end{aligned} \tag{6.15}$$

One can check that $|a_n \cos n\theta + b_n \sin n\theta| \leqslant \sqrt{a_n^2 + b_n^2}$ and then by (6.3) and Hölder's inequality:

$$\begin{aligned} d_\infty(K, S_K) &= \max_\theta \left|p_{\partial K}(\theta) - p_{\partial S_K}(\theta)\right| \\ &= \max_\theta \left| \sum_{\substack{n=3,\\ n \text{ is odd}}}^{\infty} (a_n \cos n\theta + b_n \sin n\theta) \right| \\ &\leqslant \max_\theta \left(\sum_{\substack{n=3,\\ n \text{ is odd}}}^{\infty} |a_n \cos n\theta + b_n \sin n\theta| \right) \\ &\leqslant \sum_{\substack{n=3,\\ n \text{ is odd}}}^{\infty} \frac{1}{\sqrt{n^2-1}} \cdot \sqrt{n^2-1}\sqrt{a_n^2 + b_n^2} \end{aligned}$$

$$\leqslant \sqrt{\sum_{\substack{n=3,\\ n\ is\ odd}}^{\infty} \frac{1}{n^2-1}} \cdot \sqrt{\sum_{\substack{n=3,\\ n\ is\ odd}}^{\infty} (n^2-1)(a_n^2+b_n^2)}$$
$$= \sqrt{\frac{1}{4}} \cdot \sqrt{\frac{\Phi(K)}{2\pi^2}}.$$

And the equality holds if and only if $a_{2m+1} = b_{2m+1} = 0$ for all $m \in \mathbb{N}$, so ∂K is centrally symmetric. □

Theorem 6.9. *Under the same assumptions of Theorem 6.8, one gets*

$$L_{\partial K}^2 - 4\pi A_{\partial K} - 8\left|\widetilde{A}_{CWMS(\partial K)}\right| \geqslant 16\pi d_2^2(K, S_K), \tag{6.16}$$

where equality holds if and only if ∂K is centrally symmetric or the Minkowski support function of ∂K is in the form

$$p_{\partial K}(\theta) = a_0 + a_1\cos\theta + b_1\sin\theta + a_3\cos 3\theta + b_3\sin 3\theta + \sum_{\substack{n=2,\\ n\ is\ even}}^{\infty} (a_n\cos n\theta + b_n\sin n\theta).$$

Proof. By (6.13) and (6.15)

$$\begin{aligned} d_2^2(K, S_K) &= \int_0^{2\pi} \left|p_{\partial K}(\theta) - p_{\partial S_K}(\theta)\right|^2 d\theta \\ &= \int_0^{2\pi} \left|\sum_{\substack{n=3,\\ n\ is\ odd}}^{\infty} (a_n\cos n\theta + b_n\sin\theta)\right|^2 d\theta \\ &= \pi \sum_{\substack{n=3,\\ n\ is\ odd}}^{\infty} (a_n^2+b_n^2) \\ &\leqslant \frac{1}{16\pi}\cdot 2\pi^2 \sum_{\substack{n=3,\\ n\ is\ odd}}^{\infty} (n^2-1)(a_n^2+b_n^2) \\ &= \frac{1}{16\pi}\Phi(K). \end{aligned}$$

And the equality holds if and only if $a_{2m+1} = b_{2m+1} = 0$ for all $m \in \mathbb{N}$, so ∂K is a centrally symmetric curve, or $p_{\partial K}(\theta) = a_0 + a_1\cos\theta + b_1\sin\theta + a_3\cos 3\theta + b_3\sin 3\theta + \sum_{\substack{n=2,\\ n\ \text{is even}}}^{\infty} (a_n\cos n\theta + b_n\sin n\theta)$. □

Corollary 6.10. *Under the same assumptions of Theorem 6.8, one gets*

$$\left|\widetilde{A}_{E_{\frac{1}{2}}(\partial K)}\right| \geqslant \max\left\{\pi d_\infty^2(K, S_K),\ 2d_2^2(K, S_K)\right\}, \tag{6.17}$$

(6.18)

$$\left|\widetilde{A}_{SMS(\partial K)}\right| \geqslant \max\left\{2\pi d_\infty^2(K,S_K)+\pi d_\infty^2(K,W_K),\ 2\pi d_\infty^2(K,S_K)+\frac{3}{2}d_2^2(K,W_K),\right.$$
$$\left. 4d_2^2(K,S_K)+\pi d_\infty^2(K,W_K),\ 4d_2^2(K,S_K)+\frac{3}{2}d_2^2(K,W_K)\right\},$$

where (6.18) holds whenever ∂K is not a curve of constant width.

Proof. It is an easy consequence of isoperimetric equalities (see Theorem 5.1–5.2 and Corollary 5.5) and stability properties of improved isoperimetric inequalities (see Corollary 6.5 and Theorem 6.8–6.9). □

Acknowledgements

The work of M. Zwierzyński was partially supported by NCN grant no. DEC-2013/11/B/ST1/03080. The author is sincerely grateful to Professor Wojciech Domitrz for the valuable discussions.

References

[1] Berry, M.V.: *Semi-classical mechanics in phase space: a study of Wigner's function*, Philos. Trans. R. Soc. Lond. A 287, 237-271 (1977).

[2] Barbier, E. (1860): *Note sur le probleme de l'aiguille et le jeu du joint couvert* (PDF), Journal de mathématiques pures et appliquées, 2e série (in French) 5: 273–286. See in particular pp. 283–285.

[3] Chavel, I.: *Isoperimetric Inequalities*. Differential Geometric and Analytic Perspectives. Cambridge University Press, 2001.

[4] Craizer, M.: *Iterates of Involutes of Constant Width Curves in the Mikowski Plane*, Beiträge zur Algebra und Geometrie, 55(2), 479–496, 2014.

[5] Craizer, M., Domitrz, W., Rios, P. de M.: *Even Dimensional Improper Affine Spheres* , J. Math. Anal. Appl. 421 (2015), pp. 1803–1826.

[6] DeTurck, D., Gluck, H., Pomerleano, D.: *The Four Vertex Theorem and Its Converse*, Notices Amer. Math. Soc, 54(2007), no. 2, 192–207.

[7] Domitrz, W., Janeczko, S., Rios, P. de M., Ruas, M. A. S.: *Singularities of affine equidistants: extrinsic geometry of surfaces in 4–space*, to appear in Bull. Braz. Math. Soc.

[8] Domitrz, W., Manoel, M., Rios, P. de M.: *The Wigner caustic on shell and singularities of odd functions* , Journal of Geometry and Physics 71(2013), pp. 58-72

[9] Domitrz, W., Rios, P. de M.: *Singularities of equidistants and Global Centre Symmetry sets of Lagrangian submanifolds*, Geom. Dedicata 169 (2014), pp. 361-382.

[10] Domitrz, W., Rios, P. de M., Ruas, M. A. S.: *Singularities of affine equidistants: projections and contacts*, J. Singul. 10 (2014), 67-81

[11] Domitrz, W., Zwierzyński, M.: *The geometry of the Wigner caustic and affine equidistants of planar curves*, arXiv:1605.05361

[12] Farin, G., Hoschek, J., Kim, M.-S.: *Handbook of Computer Aided Geometric Design*. North-Holland

[13] Ferone, V., Kawohl, B., Nitsch, C.: *The elastica problem under area constraint*, Mathematische Annalen, DOI 10.1007/s00208-015-1284-y

[14] Fisher, J. C.: *Curves of Constant Width from a Linear Viewpoint*, Mathematics Magazine Vol. 60, No. 3 (Jun., 1987), pp. 131-140

[15] Fuglede, B.: *Stability in the isoperimetric problem*, Bull. London Math. Soc. 18 (1986), 599 – 605

[16] Gage, M. E.: *Curve shortening makes convex curves circular*, Invent. Math., 76(1984), 357–364.

[17] Gage, M.E.: *An isoperimetric inequality with applications to curve shortening*, Duke Math. J. 50 (1983), no. 4, 1225–1229.

[18] Giblin, P.J.: *Affinely invariant symmetry sets*. Geometry and Topology of Caustics, Banach Center Publications, vol.82 (2008), p.71-84.
[19] Giblin, P.J., Warder, J.P.: *Reconstruction from medial representations*, American Mathematical Monthly 118 (2011), 712–725
[20] Giblin, P.J., Warder, J.P., Zakalyukin, V.M.: *Bifurcations of affine equidistants*, Proceedings of the Steklov Institute of Mathematics 267 (2009), 57–75.
[21] Góźdź, S.: *An antipodal set of a periodic function*, J. of Math. Anal. and App. Vol. 148, pp. 11–21, May 1990.
[22] Green, M., Osher, S.: *Steiner polynomials, Wulff flows, and some new isoperimetric inequalities for convex plane curves*, Asia J. Math., 3(1999), 659–676.
[23] Groemer, H.: *Geometric applications of Fourier series and spherical harmonics*, Encyclopedia of Mathematics and its Applications, vol. 61. Cambridge University Press, Cambridge (1996)
[24] Groemer, H.: *Stability properties of geometric inequalities*, in: P.M. Gruber, J.M. Wills (Eds.), *Handbook of Convex Geometry*, North–Holland, 1993, pp. 125–150.
[25] Grunbaum, B.: *Convex Polytopes*, John Wiley & Sons, LondonNew YorkSydney, 1967.
[26] Hsiung, C. C.: *A First Course in Differential Geometry. Pure Appl. Math.*, Wiley, New York 1981.
[27] Hurwitz, A.: *Sur quelque applications geometriques des series Fourier*, Ann. Sci. Ecole Normal Sup. (3) 19 (1902). 357–408.
[28] Hoschek, J., Lasser D.: *Fundamentals of Computer Aided Geometric Design*. A. K. Peters Wellesley MA., Ltd.
[29] Janeczko, S., Jelonek, Z., Ruas, M. A. S.: *Symmetry defect of algebraic varieties*, Asian J. Math. Vol. 18, No. 3, pp. 525-544, July 2014.
[30] Kneser, A.: *Bemerkungen uber die Anzahl der Extrema des Krummung auf geschlossenen Kurven und uber verwandte Fragen in einer nicht euklidischen Geometrie*, Festschrift Heinrich Weber. Teubner. (1912), pp. 170–180.
[31] Kimia, B., Giblin, P.J., Pollitt, A.: *Transitions of the 3D medial axis under a one-parameter family of deformations*. IEEE Transactions in Pattern Analysis and Machine Intelligence 31 (2009), 900–918.
[32] Lawlor, G.: *A new area-maximization proof for the circle*. Math. Intell. 20 (1998), 29–31
[33] Laugwitz, D.: *Differential and Riemannian Geometry*, Academic Press, New York/ London, 1965.
[34] Mukhopadhyaya, S.: *New methods in the geometry of a plane arc*, Bull. Calcutta Math. Soc. 1 (1909): 21–27.
[35] Nuño Ballesteros, J.J., Romero Fuster, M.C.: *A four vertex theorem for strictly convex space curves*, J. Geom. 46, 119126 (1993).
[36] Peternell M., Pottmann H.: *A Laguerre Geometric Approach to Rational Offsets*. Computer Aided Geometric Design vol. 15, pp. 223–249.
[37] Pan, S.L., Xu, H.: *Stability of a reverse isoperimetric inequality*. J. Math. Anal. Appl. 350 (2009), no. 1, 348-353.
[38] Reeve, G. M., Zakalyukin, V. M.: *Singularities of the Minkowski set and affine equidistants for a curve and a surface*. Topology Appl. 159 (2012), no. 2, 555–561.
[39] Steiner, J.: *Sur le maximum et le minimum des figures dans le plan, sur la sphére, et dans l'espace en général*, I and II. J. Reine Angew. Math. (Crelle) 24 (1842), 93-152 and 189-250.
[40] Schender, R.: *Convex Bodies: The Brunn-Minkowski Theory*, Cambridge, 2013.
[41] Schneider, R.: *A stability estimate for the Aleksandrov–Fenchel inequality, wich an application to mean curvature*, Manuscripta Math. 69 (1990) 291–300.
[42] Schneider, R.: *Krümmungsschwerpunkte konvexer Körper I*, Abh. Math. Sem. Univ. Hamburg 37 (1972) 112–132.
[43] Schneider, R.: *Krümmungsschwerpunkte konvexer Körper II*, Abh. Math. Sem. Univ. Hamburg 37 (1972) 204–217
[44] Thom, R.: *Structural stability and morphogenesis*. Reading, Mass: Benjamin (1975).
[45] Zakalyukin, V.M.: *Envelopes of families of wave fronts and control theory*, Proc. Steklov Math. Inst. 209 (1995), 133-142.
[46] Zwierzyński, M.: *The improved isoperimetric inequality and the Wigner caustic of planar ovals*, J. Math. Anal. Appl. 442 (2016) 726739, http://dx.doi.org/j.jmaa.2016.05.016

Warsaw University of Technology, Faculty of Mathematics and Information Science, ul. Koszykowa 75, 00-662 Warszawa, Poland
E-mail address: zwierzynskim@mini.pw.edu.pl